arXiv:1403.2236v1 [math.NT] 10 Mar 2014

ON FINITE SUMS INVOLVING $(q)_n$

MOHAMED EL BACHRAOUI

Abstract. We use an elementary argument to prove some finite sums involving expressions of the forms $(q)_n$ and $(a\,;q)_n$ along with inductive formulas for some sequences.

1. Introduction

Throughout the paper we will be using the following notation. For complex numbers q and a let

$$(a)_0=(a\,;q)_0=1,\quad (a)_n=(a\,;q)_n=\prod_{k=0}^{n-1}(1-aq^k),\quad (n\geq 1),$$

$$(a)_\infty=(a\,;q)_\infty=\prod_{k=0}^{\infty}(1-aq^k),\quad |q|<1.$$

On taking the logarithm and differentiating with respect to q in the generating function for the partition function $p(n)$

$$\prod_{n=1}^{\infty}(1-q^n)^{-1}=\sum_{n=0}^{\infty}p(n)q^n,$$

we obtain the following well-known inductive formula for the function $p(n)$

$$p(n)=\sum_{k=1}^{n}p(n-k)\sigma(k),$$

see for instance Apostol [2]. In general, if $a(n)$ is an arithmetic function with generating function

$$\prod_{n=1}^{\infty}(1-q^n)^{\frac{-f(n)}{n}}=\sum_{n=0}^{\infty}a(n)q^n,$$

then it is classical to take logarithms and to differentiate with respect to q to get inductive formulas for $a(n)$, see Apostol [2] and Robbins [5]. In El Bachraoui [4] this approach has been extended to the case

$$\prod_{i=1}^{r}\prod_{n\in A_i}(1-q^n)^{-\frac{f_i(n)}{n}}=\sum_{n=0}^{\infty}p_{\underline{A},\underline{f}}(n)q^n,$$

where $r\in\mathbb{N}$, $A_1,A_2,\dots A_\alpha\subseteq\mathbb{N}$, $\underline{A}=(A_1,A_2,\dots,A_\alpha)$, $f_i:A_i\to\mathbb{C}$ for $i=1,2,\dots,\alpha$ are arithmetic functions, and $\underline{f}=(f_1,f_2,\dots,f_\alpha)$.

1991 *Mathematics Subject Classification.* 11A25.

Key words and phrases. q-series; convolution sums; inductive formulas; arithmetic functions.

In this note we shall consider products of the form

$$\prod_{j=1}^{r}\left((1-a_jtq^{\beta_j})^{-b_j}\prod_{n=1}^{\infty}(1-a_jtq^{\alpha_jn+\beta_j})^{-\frac{f_j(n)}{n}}\right),$$

for complex constants $a_1,\dots,a_r,\ b_1,\dots,b_r$ and arithmetic functions $f_1,\dots,f_r$. However, after taking logarithms we shall differentiate with respect to t rather than q. As a result we will prove the following theorems.

Theorem 1. *If n is a positive integer and q is a complex number such that $|q|<1$, then*

$$(a)\quad \sum_{k=1}^{n}\frac{1}{1-q^k}\cdot\frac{1}{(q)_{n-k}}=n\frac{1}{(q)_n}$$

$$(b)\quad \sum_{k=1}^{n}(-1)^{k-1}\frac{1}{1-q^k}\cdot\frac{q^{\frac{(n-k)(n-k-1)}{2}}}{(q)_{n-k}}=n\frac{q^{\frac{n(n-1)}{2}}}{(q)_n}$$

Theorem 2. *If n is a positive integer and q is a complex number such that $|q|<1$, then*

$$\sum_{k=1}^{\lceil\frac{n}{2}\rceil}\left(\frac{2}{1-q^{2k-1}}\sum_{l=0}^{n-2k+1}\frac{q^{\frac{(n-2k-l+1)(n-2k-l)}{2}}}{(q)_l(q)_{n-2k-l+1}}\right)=n\sum_{k=0}^{n}\frac{q^{\frac{(n-k)(n-k-1)}{2}}}{(q)_{n-k}}.$$

Theorem 3. *Let n be a positive integer and let a and q be complex numbers such that $|q|<1$. Then*

$$\sum_{k=1}^{n}\frac{(a)_{n-k}}{(q)_{n-k}}\frac{1-a^k}{1-q^k}=n\frac{(a)_n}{(q)_n}.$$

Definition 1. [1, p. 49] The *Roger-Szego polynomial* is

$$H_n(x)=\sum_{j=0}^{n}\begin{bmatrix}n\\j\end{bmatrix}x^j,$$

where $\begin{bmatrix}n\\j\end{bmatrix}$ is the *Gaussian polynomial* given by

$$\begin{bmatrix}n\\m\end{bmatrix}=\begin{cases}\frac{(q)_n}{(q)_m(q)_{n-m}} & \text{if } 0\le m\le n,\\ 0 & \text{otherwise.}\end{cases}$$

We have the following inductive formula for $H_n(x)$.

Theorem 4. *If n is a positive integer and q is a complex number such that $|q|<1$, then*

$$\begin{aligned}n\frac{H_n(x)}{(q)_n}&=\sum_{k=1}^{n}\frac{2H_{n-k}(x)}{(q)_{n-k}(1-q^k)}\\&=\sum_{k=0}^{n-1}\frac{2H_k(x)}{(q)_k(1-q^{n-k})}.\end{aligned}$$

Definition 2. [1, p. 117] Define the sequence Λ_n by

$$\frac{(-tq^2\,;q^6)_\infty(-tq^4\,;q^6)_\infty(-tq^5\,;q^6)_\infty}{(t\,;q^6)_\infty}=\sum_{n=0}^{\infty}\frac{\Lambda_nt^n}{(q^6\,;q^6)_n}.\tag{1}$$

We note that the sequence Λ_n is related to the number $c_m(n)$ of partitions $(\lambda_1\lambda_2\ldots\lambda_s)$ of n such that $m \geq \lambda_1$, $\lambda_s \neq 1$ or 3, and for $1 \leq i \leq s$ we have $\lambda_i - \lambda_{i+1} \geq 6$ with strict inequality if $\lambda_{i+1} \equiv 0, 1$ or 3 (mod 6). Refer to Andrews[1, p. 118] for more details. The following result is an inductive formula for Λ_n.

Theorem 5. *If n is a positive integer and q is a complex number such that $|q| < 1$, then*

$$\begin{aligned} n\frac{\Lambda_n}{(q^6;q^6)_n} &= \sum_{k=1}^{n} \frac{\Lambda_{n-k}}{(q^6;q^6)_{n-k}} \frac{(-1)^{k+1}(q^{2k}+q^{4k}+q^{5k})+1}{1-q^{6k}} \\ &= \sum_{k=0}^{n-1} \frac{\Lambda_k}{(q^6;q^6)_k} \frac{(-1)^{n-k+1}(q^{2(n-k)}+q^{4(n-k)}+q^{5(n-k)}+1}{1-q^{6(n-k)}}. \end{aligned}$$

2. Main Theorem and its proof

Main Theorem. *Let $a_1,\ldots,a_r$, $b_1,\ldots,b_r$ be complex numbers, let $\alpha_1,\ldots,\alpha_r$, $\beta_1,\ldots,\beta_r$ be nonnegative integers, and let $f_1,\ldots,f_r$ be arithmetical functions such that the product*

$$F_q(t) = \prod_{j=1}^{r} \left((1-a_j t q^{\beta_j})^{-b_j} \prod_{n=1}^{\infty} (1-a_j t q^{\alpha_j n+\beta_j})^{-\frac{f_j(n)}{n}} \right)$$

and the series

$$\sum_{j=1}^{r}\sum_{n=1}^{\infty} \frac{f_j(n)}{n} q^n t^n$$

converge absolutely for $|q| < 1$ and $|t| < 1$ and represent analytic functions in the unit disk $|t| < 1$ for $|q| < 1$. If $F_q(t) = \sum_{n=0}^{\infty} A_q(n)t^n$, then $A_q(0) = 1$ and for all $n \geq 1$ we have

$$nA_q(n) = \sum_{k=0}^{n-1} A_q(k)h(n-k) = \sum_{k=1}^{n} A_q(n-k)h(k), \tag{2}$$

where

$$h(m) = \sum_{j=1}^{r} b_j a_j^m q^{\beta_j m} + \sum_{j=1}^{r}\sum_{k=1}^{\infty} a_j^m \frac{f(k)}{k} q^{m(k\alpha_j+\beta_j)}.$$

Proof. It is clear that $A_q(0) = 1$. Now suppose that $n \geq 1$. It is easily seen that the second identity of formula (2) holds. Further we have

$$\begin{aligned} \log F_q(t) &= \sum_{j=1}^{r} -b_j \log(1-a_j t q^{\beta_j}) + \sum_{j=1}^{r}\sum_{n=1}^{\infty} -\frac{f_j(n)}{n} \log(1-a_j t q^{\alpha_j n+\beta_j}) \\ &= \sum_{j=1}^{r} b_j \sum_{m=1}^{\infty} \frac{a_j^m q^{\beta_j m}}{m} t^m + \sum_{j=1}^{r}\sum_{n=1}^{\infty} \frac{f_j(n)}{n} \sum_{m=1}^{\infty} \frac{a_j^m q^{m(\alpha_j n+\beta_j)}}{m} t^m, \end{aligned}$$

from which it follows that

$$t\frac{\frac{d}{dt}F_q(t)}{F_q(t)} = t\frac{d}{dt}\log F_q(t) = \sum_{m=1}^{\infty}\sum_{j=1}^{r} \left(b_j a_j^m q^{\beta_j m} + \sum_{n=1}^{\infty} \frac{f_j(n)}{n} a_j^m q^{m(\alpha_j n+\beta_j)} \right) t^m = \sum_{m=1}^{\infty} h(m)t^m,$$

where

$$h(m)=\sum_{j=1}^{r}b_j a_j^m q^{\beta_j m}+\sum_{j=1}^{r}\left(\sum_{\substack{m|k\\ \frac{k-\beta_j m}{\alpha_j m}\in\mathbb{N}}}\frac{a_j^m f_j(\frac{k-\beta_j m}{\alpha_j m})}{\frac{k-\beta_j m}{\alpha_j m}}\right)q^k$$
$$=\sum_{j=1}^{r}b_j a_j^m q^{\beta_j m}+\sum_{j=1}^{r}\sum_{k=1}^{\infty}a_j^m\frac{f(k)}{k}q^{m(k\alpha_j+\beta_j)}.$$

Thus

$$t\frac{d}{dt}F_q(t)=F_q(t)\sum_{m=0}^{\infty}h(m)t^m,$$

or equivalently,

$$\sum_{n=1}^{\infty}nA_q(n)t^n=\sum_{n=0}^{\infty}A_q(n)t^n\cdot\sum_{n=0}^{\infty}h(n)t^n,$$

which after equating coefficients gives the desired identity. □

3. Proofs

We note that the well-know facts used in the proofs are all found in Andrews [1] or Berndt [3].

Proof of Theorem 1. (a) A well-known result of Euler states that

$$\prod_{n=0}^{\infty}(1-tq^n)^{-1}=1+\sum_{n=1}^{\infty}\frac{t^n}{(q)_n},$$

or equivalently,

$$(1-t)^{-1}\prod_{n=1}^{\infty}(1-tq^n)^{-1}=\sum_{n=0}^{\infty}\frac{t^n}{(q)_n}. \tag{3}$$

Then an application of Main Theorem to $r=1,\ a_1=1,\ b_1=1,\ \alpha_1=1,\ \beta_1=0,$ and $f_1(n)=n$ gives

$$h(m)=1+\sum_{m|k}q^k=\frac{1}{1-q^m}$$

and therefore by identity (2) we have for all $n\geq 1$ that

$$\frac{n}{(q)_n}=\sum_{k=1}^{n}\frac{1}{1-q^k}\cdot\frac{1}{(q)_{n-k}},$$

as desired.

(b) By another result of Euler we have

$$\prod_{n=0}^{\infty}(1+tq^n)=1+\sum_{n=1}^{\infty}\frac{q^{\frac{n(n-1)}{2}}t^n}{(q)_n},$$

or equivalently,

$$(1+t)\prod_{n=1}^{\infty}(1+tq^n)=\sum_{n=0}^{\infty}\frac{q^{\frac{n(n-1)}{2}}t^n}{(q)_n}. \tag{4}$$

Then by Main Theorem applied to $r=1,\ a_1=-1,\ b_1=-1,\ \alpha_1=1,\ \beta_1=0$, and $f_1(n)=-n$ we get

$$h(m)=(-1)^{m+1}+(-1)^{m+1}\sum_{m|k}q^k=\frac{(-1)^{m+1}}{1-q^m}$$

and thus by formula (2)

$$n\frac{q^{\frac{n(n-1)}{2}}}{(q)_n}=\sum_{k=1}^{n}(-1)^{k-1}\frac{1}{1-q^k}\cdot\frac{q^{\frac{(n-k)(n-k-1)}{2}}}{(q)_{n-k}},$$

which completes the proof.

Proof of Theorem 2. By Euler's formulas (3) and (4) we have

$$\begin{aligned}(1-t)^{-1}(1+t)\prod_{n=1}^{\infty}(1-tq^n)^{-1}(1+tq^n)&=\sum_{n=0}^{\infty}\frac{t^n}{(q)_n}\cdot\sum_{n=0}^{\infty}\frac{q^{\frac{n(n-1)}{2}}t^n}{(q)_n}\\&=\sum_{n=0}^{\infty}\left(\sum_{k=0}^{n}\frac{q^{\frac{(n-k)(n-k-1)}{2}}}{(q)_{n-k}}\right)t^n.\end{aligned}$$

Now apply Main Theorem to the last identity with $r=2,\ a_1=1,\ b_1=1,\ \alpha_1=1,\ \beta_1=0,\ f_1(n)=n,\ a_2=-1,\ b_2=-1,\ \alpha_2=1,\ \beta_2=0$, and $f_2(n)=-n$ to obtain

$$h(m)=1+(-1)^{m-1}+\sum_{m|k}(1+(-1)^{m-1})q^k=\begin{cases}\frac{2}{1-q^m}, & \text{if } m \text{ is odd}\\ 0, & \text{otherwise}\end{cases}$$

and to deduce the desired formula from identity (2).

Proof of Theorem 3. By a well-known result of Cauchy, also referred to as the q-analogue of the binomial theorem, we have for $|q|<1$ and $|t|<1$

$$\prod_{n=0}^{\infty}\frac{1-atq^n}{1-tq^n}=1+\sum_{n=1}^{\infty}\frac{(a)_n}{(q)_n}t^n,$$

or equivalently,

$$(1-at)(1-t)^{-1}\prod_{n=1}^{\infty}(1-atq^n)(1-tq^n)^{-1}=\sum_{n=0}^{\infty}\frac{(a)_n}{(q)_n}t^n.$$

Letting in Main Theorem, $r=2,\ a_1=a,\ a_2=1,\ b_1=-1,\ b_2=1,\ \alpha_1=\alpha_2=1,\ \beta_1=\beta_2=0,\ f_1(n)=-n$, and $f_2(n)=n$, we obtain

$$h(m)=(1-a^m)+\sum_{m|k}(1-a^m)q^k=\frac{1-a^m}{1-q^m}$$

and therefore

$$n\frac{(a)_n}{(q)_n}=\sum_{k=1}^{n}\frac{(a)_k}{(q)_k}\frac{1-a^{n-k}}{1-q^{n-k}},$$

which completes the proof.

Proof of Theorem 4. It is well-known that

$$\frac{1}{(t)_\infty(tx)_\infty}=\sum_{n=0}^{\infty}\frac{H_n(x)}{(q)_n}t^n,$$

or equivalently,

$$(1-t)^{-1}(1-xt)^{-1}\prod_{n=1}^{\infty}(1-tq^n)^{-1}(1-xtq^n)^{-1}=\sum_{n=0}^{\infty}\frac{H_n(x)}{(q)_n}t^n.$$

Now apply Main Theorem to $r=2,\ a_1=b_1=1,\ \alpha_1=1,\ \beta_1=0,\ f_1(n)=n,$ $a_2=x,\ b_2=1,\ \ \alpha_2=1,\ \beta_2=0$, and $f_2(n)=n$ to get

$$h(m)=2+\sum_{m|k}2q^k=\frac{2}{1-q^m},$$

and to derive the desired identity.

Proof of Theorem 5. The left hand side of identity (1) is clearly equal to

$$(1+tq^2)(1+tq^4)(1+tq^5)(1-t)^{-1}\prod_{n=1}^{\infty}(1+tq^{6n+2})(1+tq^{6n+4})(1+tq^{6n+5})(1-tq^{6n})^{-1}.$$

Appealing to Main Theorem with $a_1=a_2=a_3=-1,\ a_4=1,\ b_1=b_2=b_3=-1,\ b_4=1,\ \alpha_1=\alpha_2=\alpha_3=\alpha_4=6,\ \beta_1=2,\ \beta_2=4,\ \beta_3=5,\ \beta_4=0,$ $f_1(n)=f_2(n)=f_3(n)=-n$, and $f_4(n)=n$ yields

$$\begin{aligned}h(m)&=(-1)^{m+1}(q^{2m}+q^{4m}+q^{5m})+\frac{(-1)^{m+1}}{1-q^{6m}}(q^{8m}+q^{10m}+q^{11m})+\frac{1}{1-q^{6m}}\\&=\frac{1}{1-q^{6m}}((-1)^{m+1}(q^{2m}+q^{4m}+q^{5m})+1),\end{aligned}$$

which if put in formula (2) gives the desired identity.

References

[1] G. E. Andrews, *The theory of partitions*, Cambridge University Press, 1998.

[2] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer, 1976.

[3] B. C. Berndt, *Number theory in the spirit of Ramanujan*, American Matematical Society, 2006.

[4] M. El Bachraoui, *Inductive Formulas for some Arithmetic Functions*, International Journal of Number Theory 7, Number 8, (2011), 2261-2268.

[5] N. Robbins, *Some identities connecting partition functions to other number theoretic functions*, Rocky Mountain Journal of Mathematics, Volume 29, Number 1, (1999), 335-345.

Dept. Math. Sci, United Arab Emirates University, PO Box 17551, Al-Ain, UAE

E-mail address: melbachraoui@uaeu.ac.ae